

TAYLOR SPECTRUM AND CHARACTERISTIC FUNCTIONS OF COMMUTING 2-CONTRACTIONS

BERRABAH BENDOUKHA

ABSTRACT. In this paper, we give a description of Taylor spectrum of commuting 2-contractions in terms of characteristic functions of such contractions. The case of a single contraction obtained by B. Sz. Nagy and C. Foias is generalised in this work.

1. INTRODUCTION

Let $\mathcal{H}$ be a Hilbert space. A 2-tuple $A = (A_1, A_2)$ of bounded operators on $\mathcal{H}$ is called contractive (or 2-contraction) if $\|A_1 h_1 + A_2 h_2\|^2 \leq \|h_1\|^2 + \|h_2\|^2$ for all h_1, h_2 in $\mathcal{H}$. It is equivalent [1] to the condition: $A_1 A_1^* + A_2 A_2^* \leq 1_{\mathcal{H}}$. If additionally, operators A_1 and A_2 commute, then A is called a commuting 2-contraction. To every commuting 2-contraction $A = (A_1, A_2)$ corresponds an analytic operator-valued function $\theta_A : \mathbb{D}^2 \rightarrow B(\mathcal{D}_A, \mathcal{D}_{A^*})$ called characteristic function of A and defined by :

$$(1.1) \quad \theta_A(z_1, z_2) = -A + D_{A^*} (1_{\mathcal{H}} - z_1 A_1^* - z_2 A_2^*)^{-1} (z_1 \cdot 1_{\mathcal{H}}, z_2 \cdot 1_{\mathcal{H}}) D_A$$

where

- (a): $\mathbb{D}^2 = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 < 1\}$,
- (b): $B(\mathcal{D}_A, \mathcal{D}_{A^*})$ is the set of all bounded operators from $\mathcal{D}_A$ into $\mathcal{D}_{A^*}$,
- (c): $D_{A^*} = (I - A_1 A_1^* - A_2 A_2^*)^{\frac{1}{2}} : \mathcal{H} \rightarrow \mathcal{H}$ and $\mathcal{D}_{A^*}$ is the closure of the range of D_{A^*} ,
- (d): $D_A = \begin{bmatrix} 1_{\mathcal{H}} - A_1^* A_1 & -A_1^* A_2 \\ -A_2^* A_1 & 1_{\mathcal{H}} - A_2^* A_2 \end{bmatrix}^{\frac{1}{2}} : \mathcal{H}^2 \rightarrow \mathcal{H}^2$ and is the closure of the range of D_A ,
- (e): $(z_1 1_{\mathcal{H}}, z_2 1_{\mathcal{H}}) : \mathcal{H}^2 \rightarrow \mathcal{H}$; $(z_1 1_{\mathcal{H}}, z_2 1_{\mathcal{H}}) \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = z_1 h_1 + z_2 h_2$.

Characteristic function of commuting n -contraction has been introduced in [3] as a generalization of characteristic function of a single contraction [9]. A lot of its remarkable properties have been established in ([2], [4],[9]). In particular, it is shown (like in the single case) that the characteristic function is a unitary invariant. It means that characteristic functions of two pure or completely noncoisometric n -contractions ([2], [4]) $A = (A_1, \dots, A_n)$ and $A' = (A'_1, \dots, A'_n)$ coincide if and only

2000 *Mathematics Subject Classification.* 47A10, 47A13.

Key words and phrases. Characteristic function, 2-contraction, Taylor spectrum.

if there exists a unitary operator $U : \mathcal{H} \rightarrow \mathcal{H}$ such that $A'_i = U^{-1}A_iU$ for every $i = 1, \dots, n$.

In the case of a completely nonunitary single contraction, the spectrum can be described in terms of the characteristic function (see theorem 4.1 [9]). The aim of this paper is to give a description of Taylor spectrum in the case of commuting pure 2-contractions by means of characteristic function (1.1).

In section 2 we briefly remind the definition of Taylor spectrum. Section 3 contains characterizations of different components of Taylor spectrum. In section 4, we investigate the behavior of Taylor spectrum under the action of involutive automorphisms of unit ball.

2. TAYLOR SPECTRUM

Let $A = (A_1, A_2, \dots, A_n)$ be a pure n -contraction. According to ([5], [6], [7], [10]), the Taylor spectrum of A can be defined as follows. Let $\Lambda(\mathcal{H})$ be the exterior algebra on n generators $e_1, \dots, e_n$ with identity $e_0 = 1$ and coefficients in H . In other words,

$$\Lambda(\mathcal{H}) = \{x \otimes e_{i_1} \wedge \dots \wedge e_{i_p} : x \in \mathcal{H}; 1 \leq i_1, \dots, i_p \leq n; 1 \leq p \leq n\}$$

with the collapsing property : $e_i \wedge e_j + e_j \wedge e_i = 0$. One has

$$\begin{aligned} \Lambda(\mathcal{H}) &= \bigoplus_{k=1}^n \Lambda^k(\mathcal{H}); & \Lambda^k(\mathcal{H}) &= \{x \otimes e_{i_1} \wedge \dots \wedge e_{i_k} : x \in H, 1 \leq i_1, \dots, i_k \leq n\}; \\ \Lambda^0(\mathcal{H}) &= \mathcal{H}. \end{aligned}$$

Consider in $\Lambda(\mathcal{H})$ operator:

$$B_A : \Lambda(\mathcal{H}) \rightarrow \Lambda(\mathcal{H}) : \quad B_A(x \otimes e_{i_1} \wedge \dots \wedge e_{i_p}) = \sum_{k=1}^n A_k(x) \otimes e_k \wedge e_{i_1} \wedge \dots \wedge e_{i_p}.$$

It is not difficult to see that $B_A^2 = 0$ and $\text{Ran } B_A \subseteq \text{Ker } B_A$. Decomposition $\Lambda(\mathcal{H}) = \bigoplus_{k=1}^n \Lambda^k(\mathcal{H})$ gives a rise to a cochain $K(A, \mathcal{H})$, the so-called Koszul complex $K(A, \mathcal{H})$ associated to A on $\mathcal{H}$ as follows:

$$K(A_1, A_2, \mathcal{H}) : \{0\} \rightarrow \mathcal{H} = \Lambda^0(\mathcal{H}) \xrightarrow{B_A^0} \dots \xrightarrow{B_A^{n-1}} \Lambda^n(\mathcal{H}) \rightarrow \{0\}$$

where B_A^k is the restriction of B_A to the subspace $\Lambda^k(\mathcal{H})$. Complex $K(A, \mathcal{H})$ is said to be exact (or regular) if:

$$\{0\} = \ker B_A^0, \quad \text{Ran } B_A^0 = \ker B_A^1, \dots, \quad \text{Ran } B_A^{n-2} = \ker B_A^{n-1}, \quad \text{Ran } B_A^{n-1} = \Lambda^n(\mathcal{H}).$$

Definition 1. *The Taylor spectrum of n -contraction is the set:*

$$\sigma_T(A) = \{z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n : K(A_1 - z_1, \dots, A_n - z_n; H) \text{ is not exact}\}.$$

Let us now suppose that $n = 2$. Then,

$$\begin{aligned} \Lambda(\mathcal{H}) &= \Lambda^0(\mathcal{H}) \oplus \Lambda^1(\mathcal{H}) \oplus \Lambda^2(\mathcal{H}) \\ &= (\mathcal{H} \otimes e_0) \oplus ((\mathcal{H} \otimes e_1) \oplus (\mathcal{H} \otimes e_2)) \oplus (\mathcal{H} \otimes e_1 \wedge e_2). \end{aligned}$$

According to this direct sum, operator B_A admits the matrix representation

$$B_A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ A_1 & 0 & 0 & 0 \\ A_2 & 0 & 0 & 0 \\ 0 & -A_2 & A_1 & 0 \end{bmatrix}$$

and then operators $B_{(A_1, A_2)}^0$ and $B_{(A_1, A_2)}^1$ have the forms:

$$(2.1) \quad \begin{cases} B_{(A_1, A_2)}^0(x) = A_1(x) \oplus A_2(x), & (x, y \in \mathcal{H}), \\ B_{(A_1, A_2)}^1(x \oplus y) = -A_2(x) + A_1(y), & (x, y \in \mathcal{H}) \end{cases}.$$

According to definition 1 and formula (2.1), one has

$$\sigma_T(A_1, A_2) = \sigma_T^{(1)}(A_1, A_2) \cup \sigma_T^{(2)}(A_1, A_2) \cup \sigma_T^{(3)}(A_1, A_2)$$

where

$$(2.2) \quad (z_1, z_2) \in \sigma_T^{(1)}(A_1, A_2) \Leftrightarrow \exists x \in \mathcal{H} : x \neq 0, (A_1 - z_1)x = (A_2 - z_2)x = 0$$

$$(2.3)$$

$$(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2) \Leftrightarrow \exists (x_1, x_2) \in \mathcal{H}^2 : \begin{cases} (A_1 - z_1)x_1 - (A_2 - z_2)x_2 = 0 \\ (x_1, x_2) \neq ((A_1 - z_1)h, (A_2 - z_2)h), \forall h \in \mathcal{H} \end{cases}$$

$$(2.4)$$

$$(z_1, z_2) \in \sigma_T^{(3)}(A_1, A_2) \Leftrightarrow \exists y \in \mathcal{H} : y \neq (A_1 - z_1)x_1 - (A_2 - z_2)x_2, \forall (x_1, x_2) \in \mathcal{H}^2.$$

Remark 1. $\sigma_T^{(1)}(A_1, A_2)$ is called the *ponctual joint Taylor spectrum*.

Taylor joint spectrum generalizes the one variable notion of spectrum. It is a nonempty compact subset of $\mathbb{C}^n$. The reader can find an excellent account of the Taylor spectrum and its relations with other multiparameter spectral theories in [6]. Note also that in [2] a description of Harte spectrum by means of characteristic function is given.

Throughout this paper, we will suppose that if $A = (A_1, A_2)$ is a commuting 2-contraction, then operator $D_{A^*} = (I - A_1 A_1^* - A_2 A_2^*)^{\frac{1}{2}}$ is one to one. Note that pure (and more generally completely non coisometric) commutig 2-contractions ([2], [3], [4]) satisfy this condition. Indeed, if $A = (A_1, A_2)$ is a pure 2-contraction then, the decreasing sequence of positive bounded operators $((A_1 A_1^* + A_2 A_2^*)^n)_{n \in \mathbb{N}}$ admits a strong limit $A_\infty = 0$. Because of that,

$$\begin{aligned} D_{A^*}(x) &= 0 \Rightarrow D_{A^*}^2(x) = (I - A_1 A_1^* - A_2 A_2^*)x = 0 \\ &\Rightarrow x = (A_1 A_1^* + A_2 A_2^*)x \Rightarrow x = (A_1 A_1^* + A_2 A_2^*)^n x, \forall n = 0, 1, 2, \dots \\ &\Rightarrow x = A_\infty(x) = 0. \end{aligned}$$

Using relations $AD_A = D_{A^*}A$ and $A^*D_{A^*} = D_A A^*$ ([3]), it can be proven that :

$$D_{A^*} = (I - A_1 A_1^* - A_2 A_2^*)^{\frac{1}{2}} \text{ is one to one } \Leftrightarrow D_A = \begin{bmatrix} 1_H - A_1^* A_1 & -A_1^* A_2 \\ -A_2^* A_1 & 1_H - A_2^* A_2 \end{bmatrix}^{\frac{1}{2}}$$

is one to one.

3. Characterization of Taylor spectrum

Lemma 1. *Let $A = (A_1, A_2)$ be a commuting 2-contraction such that D_{A^*} is one to one. Then,*

$$A(x, y) = z_1 x + z_2 y, ((z_1, z_2) \in \mathbb{C}^2, (x, y) \in \mathcal{H}^2) \Leftrightarrow \theta_A(z_1, z_2) \left(D_A \begin{pmatrix} x \\ y \end{pmatrix} \right) = 0.$$

Proof. It follows directly from relation

(3.1)

$$\theta_A(z_1, z_2) D_A \begin{pmatrix} x \\ y \end{pmatrix} = D_{A^*} (1_{\mathcal{H}} - z_1 A_1^* - z_2 A_2^*)^{-1} [(z_1 x + z_2 y) - (A_1 x + A_2 y)]$$

□

Lemma 2. *Let $A = (A_1, A_2)$ be a commuting pure 2-contraction such that D_{A^*} is one to one, $(z_1, z_2) \in \mathbb{C}^2$ and $x \in \mathcal{H}$. Then,*

$$A^*(x) = \begin{pmatrix} \overline{z_1} \cdot x \\ \overline{z_2} \cdot x \end{pmatrix} \Leftrightarrow (\theta_A(z_1, z_2))^* D_{A^*}(x) = 0.$$

Proof. Since in this case $D_{A^*}^2(x) = (1_{\mathcal{H}} - \overline{z_1} A_1 - \overline{z_2} A_2)x$, then the necessary condition is a direct consequence of relation,

(3.2)

$$(\theta_A(z_1, z_2))^* D_{A^*}(x) = D_A \left(- \begin{pmatrix} A_1^* x \\ A_2^* x \end{pmatrix} + \begin{pmatrix} \overline{z_1} \\ \overline{z_2} \end{pmatrix} \right) \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) D_{A^*}^2(x)$$

Proof of the suffisant condition Hence $(\theta_A(0, 0))^* D_{A^*}(x) = - \begin{pmatrix} A_1^* x \\ A_2^* x \end{pmatrix}$, one

can whithout loosing the generality suppose that $(z_1, z_2) \neq (0, 0)$.

$$\begin{aligned} (\theta_A(z_1, z_2))^* D_{A^*}(x) &= 0 \\ \Rightarrow D_A \left(- \begin{pmatrix} A_1^* x \\ A_2^* x \end{pmatrix} + \begin{pmatrix} \overline{z_1} \\ \overline{z_2} \end{pmatrix} \right) \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) D_{A^*}^2(x) &= 0 \\ \Rightarrow \begin{cases} -A_1^* x + \overline{z_1} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \\ -A_2^* x + \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \end{cases} \\ \Rightarrow \begin{cases} -A_1^* x + \overline{z_1} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \\ -A_2^* x + \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \end{cases} \\ \Rightarrow \begin{cases} -\overline{z_2} A_1^* x + \overline{z_2} \overline{z_1} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \\ -\overline{z_1} A_2^* x + \overline{z_1} \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \end{cases} \\ \Rightarrow \begin{cases} \overline{z_1} A_2^* x = \overline{z_2} A_1^* x \\ -\overline{z_2} A_1^* x + \overline{z_2} \overline{z_1} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0 \end{cases} \\ \Rightarrow \begin{cases} \overline{z_1} A_2^* x = \overline{z_2} A_1^* x \\ -\overline{z_2} A_1^* x + \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (\overline{z_1} I - \overline{z_1} A_1 A_1^* - \overline{z_2} A_2 A_1^*) x = 0 \end{cases} \\ \Rightarrow \begin{cases} \overline{z_1} A_2^* x = \overline{z_2} A_1^* x \\ \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) ((\overline{z_1} I - A_1^*)) x = 0 \end{cases} \\ \Rightarrow ((\overline{z_1} I - A_1^*)) x = 0 \Rightarrow A_1^* x = \overline{z_1} x. \end{aligned}$$

On the other hand,

- (1) $\overline{z_1} A_2^* x = \overline{z_2} A_1^* x$, $A_1^* x = \overline{z_1} x$ and $z_1 \neq 0 \Rightarrow A_2^* x = \overline{z_2} x$.
- (2) Putting $A_1^* x = \overline{z_1} x$ and $z_1 = 0$ in the relation

$$-A_2^* x + \overline{z_2} \left((1_H - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1} \right) (I - A_1 A_1^* - A_2 A_2^*) x = 0,$$

and multiplying by $(1_H - \overline{z_2}A_2)^{-1}$, one obtains

$$(1_H - \overline{z_2}A_2) A_2^* x = \overline{z_2} (I - A_2 A_2^*) x.$$

This last relation is equivalent to $A_2^* x = \overline{z_2} x$.

□

Proposition 1. *Let $A = (A_1, A_2)$ be a commuting pure 2-contraction such that D_{A^*} is one to one. Then, $(z_1, z_2) \in \sigma_T^{(1)}(A_1, A_2)$ if and only if equation $\theta_A(z_1, z_2)X = 0$ admits at least two nontrivial solutions $D_A(X_1)$ and $D_A(X_2)$ such that, $X_1 = \begin{pmatrix} x \\ 0 \end{pmatrix}$, $X_2 = \begin{pmatrix} 0 \\ x \end{pmatrix}$, $x \in \mathcal{H}$.*

Proof. One has

$$\begin{aligned} (z_1, z_2) &\in \sigma_T^{(1)}(A_1, A_2) \\ &\Leftrightarrow \exists x \in \mathcal{H} : x \neq 0, (A_1 - z_1)x = (A_2 - z_2)x = 0 \\ &\Leftrightarrow \exists x \in \mathcal{H} : x \neq 0, A_1(x) = z_1.x \text{ and } A_2(x) = z_2.x \\ &\Leftrightarrow \exists x \in \mathcal{H} : x \neq 0, A \begin{pmatrix} x \\ 0 \end{pmatrix} = z_1.x + z_2.0 \text{ and } A \begin{pmatrix} 0 \\ x \end{pmatrix} = z_1.0 + z_2.x \\ &\Leftrightarrow \theta_A(z_1, z_2) \left(D_A \begin{pmatrix} x \\ 0 \end{pmatrix} \right) = 0 \text{ and } \theta_A(z_1, z_2) \left(D_A \begin{pmatrix} 0 \\ x \end{pmatrix} \right) = 0 \end{aligned}$$

To end the proof, it is sufficient to remark that

$$x \neq 0 \Leftrightarrow D_A \begin{pmatrix} x \\ 0 \end{pmatrix} \neq 0 \Leftrightarrow D_A \begin{pmatrix} 0 \\ x \end{pmatrix} \neq 0.$$

□

Proposition 2. *Let $A = (A_1, A_2)$ be a commuting pure 2-contraction such that D_{A^*} is one to one. Then, $(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2)$ if and only if the equation $\theta_A(z_1, z_2)Y = 0$ admits at least one non trivial solution $Y = D_A(X)$ such that $X \neq \begin{pmatrix} (A_1 - z_1)h \\ (z_2 - A_2)h \end{pmatrix}$, $\forall h \in \mathcal{H}$.*

Proof. One has,

$$(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2) \Leftrightarrow \begin{cases} \exists (x_1, x_2) \in \mathcal{H}^2 : (A_1 - z_1)x_1 - (A_2 - z_2)x_2 = 0, \\ (x_1, x_2) \neq ((A_1 - z_1)h, (A_2 - z_2)h) \forall h \in \mathcal{H}, \\ (x_1, x_2) \neq (0, 0). \end{cases}$$

It means that

$$(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2) \Leftrightarrow \begin{cases} \exists (x_1, x_2) \in \mathcal{H}^2 : A_1(x_1) + A_2(-x_2) = z_1.x_1 + z_2.(-x_2), \\ (x_1, x_2) \neq ((A_1 - z_1)h, (A_2 - z_2)h); \forall h \in \mathcal{H}, \\ (x_1, x_2) \neq (0, 0). \end{cases}$$

According to lemma 1, one has finally

$$(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2) \Leftrightarrow \begin{cases} \exists (x_1, x_2) \in \mathcal{H}^2 : \theta_A(z_1, z_2) \left(D_A \begin{pmatrix} x_1 \\ -x_2 \end{pmatrix} \right) = 0, \\ (x_1, x_2) \neq ((A_1 - z_1)h, (A_2 - z_2)h); \forall h \in \mathcal{H}, \\ (x_1, x_2) \neq (0, 0). \end{cases} \quad .$$

□

Proposition 3. *Let $A = (A_1, A_2)$ be a commuting 2-contraction such that D_{A^*} is one to one and $(z_1, z_2) \in \mathbb{C}^2$. Assume that equation $(\theta_A(z_1, z_2))^* D_{A^*}(y) = 0$ admits at least one non trivial solution. Then,*

$$(z_1, z_2) \in \sigma_T^{(3)}(A_1, A_2).$$

Proof. Suppose that $(\theta_A(z_1, z_2))^* D_{A^*}(y) = 0$ admits at least one non trivial solution y . According to lemma 2, it means that

$$A^*(y) = \begin{pmatrix} \overline{z_1} \cdot y \\ \overline{z_2} \cdot y \end{pmatrix}$$

Thus, for every $(x_1, x_2) \in \mathcal{H}^2$, one has

$$\begin{aligned} 0 &= \left\langle A^*(y) - \begin{pmatrix} \overline{z_1} \cdot y \\ \overline{z_2} \cdot y \end{pmatrix}, \begin{pmatrix} x_2 \\ -x_1 \end{pmatrix} \right\rangle_{\mathcal{H}^2} \\ &= \left\langle \begin{pmatrix} (A_1^* - \overline{z_1}) \cdot y \\ (A_2^* - \overline{z_2}) \cdot y \end{pmatrix}, \begin{pmatrix} x_2 \\ -x_1 \end{pmatrix} \right\rangle_{\mathcal{H}^2} \\ &= \langle (A_1^* - \overline{z_1}) \cdot y, x_2 \rangle_{\mathcal{H}} + \langle (A_2^* - \overline{z_2}) \cdot y, -x_1 \rangle_{\mathcal{H}} \\ &= \langle y, (A_1 - z_1)x_2 \rangle_{\mathcal{H}} - \langle y, (A_2 - z_2)x_1 \rangle_{\mathcal{H}} \\ &= \langle y, (A_1 - z_1)x_2 - (A_2 - z_2)x_1 \rangle_{\mathcal{H}}. \end{aligned}$$

Thus,

$$y \notin \text{Ran} D_{(A_1 - z_1, A_2 - z_2)}^1$$

and finally

$$(z_1, z_2) \in \sigma_T^{(3)}(A_1, A_2).$$

□

4. TAYLOR SPECTRUM AND INVOLUTIVE AUTOMORPHISMS OF UNIT BALL

In this section we investigate the Taylor spectrum under action of involutive automorphisms of unit ball $\mathbb{D}^2$. Such automorphisms are defined in [8] by

$$\Phi_\lambda(z) = \lambda - \frac{\sqrt{1 - \|\lambda\|^2}}{1 - \langle z, \lambda \rangle} \left(z - \left(1 - \sqrt{1 - \|\lambda\|^2} \right) \frac{\langle z, \lambda \rangle}{\|\lambda\|^2} \cdot \lambda \right), \lambda \in \mathbb{D}^2, \lambda = (\lambda_1, \lambda_2) \neq 0.$$

Connection between automorphisms of unit ball and multicontractions has been made in [2] and [4] where some very interesting properties have been established. In particular, if Φ_λ is an involutive automorphism of unit ball and $A = (A_1, A_2)$ is a commutative 2-contraction then, (see [2], sections 4 and 5) one can define operator

$$\Phi_\lambda(A) = \Lambda - D_{\Lambda^*} (1_H - \Lambda \Lambda^*)^{-1} \Lambda D_\Lambda$$

where the operator $\Lambda = (\lambda_1 \cdot 1_H, \lambda_2 \cdot 1_H)$ is defined from H^2 into $\mathcal{H}$ by :

$$\Lambda(x_1, x_2) = \lambda_1 \cdot x_1 + \lambda_2 \cdot x_2.$$

Proposition 4 and theorem 4.2 below summarise important for us results obtained in [2] and [4].

Proposition 4. *Let Φ_λ an involutive automorphism of unit ball and $A = (A_1, A_2)$ a commutative 2-contraction. Then, $\Phi_\lambda(A)$ is a commutative 2-contraction such that,*

$$(4.1) \quad I - \Phi_\lambda(A)^* \Phi_\lambda(A) = D_\Lambda (1_{\mathcal{H}} - A^* \Lambda)^{-1} (I - A^* A) (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda,$$

$$(4.2) \quad I - \Phi_\lambda(A) \Phi_\lambda(A)^* = D_{\Lambda^*} (1_{\mathcal{H}} - \Lambda \Lambda^*)^{-1} (I - A A^*) (1_{\mathcal{H}} - \Lambda A^*)^{-1} D_{\Lambda^*}.$$

Theorem 1. *Let Φ_λ be an involutive automorphism of unit ball and $A = (A_1, A_2)$ a commutative 2-contraction. Then,*

(1) *Operators $\Omega : D_{\Phi_\lambda(A)} \rightarrow D_A$ and $\Omega_* : D_{\Phi_\lambda(A)^*} \rightarrow D_{A^*}$ defined by*

$$\begin{aligned} \Omega(D_{\Phi_\lambda(A)}(X)) &= D_A (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda(X), \\ \Omega_*(D_{\Phi_\lambda(A)^*}(X)) &= D_{A^*} (1_{\mathcal{H}} - \Lambda A^*)^{-1} D_{\Lambda^*}(X) \end{aligned}$$

are unitaries.

(2) *$\theta_{\Phi_\lambda(A)}$ and θ_A are connected by the relation*

$$\Omega_* \theta_{\Phi_\lambda(A)}(z_1, z_2) = \theta_A(\Phi_\lambda(z_1, z_2)) \Omega.$$

It can be shown that:

$$(4.3) \quad (1_{\mathcal{H}} - \Lambda \Lambda^*)^{-1} = (1_{\mathcal{H}} - \overline{z_1} A_1 - \overline{z_2} A_2)^{-1},$$

$$(4.4) \quad (1_{\mathcal{H}^2} - \Lambda^* A)^{-1} = \begin{bmatrix} (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} (1_{\mathcal{H}} - \overline{\lambda_2} A_2) & \overline{\lambda_2} A_1 \cdot (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} \\ \overline{\lambda_1} A_2 \cdot (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} & (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} (1_{\mathcal{H}} - \overline{\lambda_1} A_1) \end{bmatrix}$$

$$(4.5) \quad D_\Lambda = \frac{1}{|\lambda_1|^2 + |\lambda_2|^2} \begin{bmatrix} |\lambda_2|^2 + |\lambda_1|^2 \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} & \overline{\lambda_1} \lambda_2 \left(\sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} - 1 \right) \\ \lambda_1 \overline{\lambda_2} \left(\sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} - 1 \right) & |\lambda_1|^2 + |\lambda_2|^2 \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} \end{bmatrix}.$$

Using (4.3), one can show that

$$(4.6) \quad \Phi_\lambda(A) = (B_1(\lambda), B_2(\lambda))$$

where

$$\begin{aligned} B_1(\lambda) &= \lambda_1 \cdot 1_{\mathcal{H}} - \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} \\ &\quad \times \left\{ A_1 \left(|\lambda_2|^2 + |\lambda_1|^2 \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} \right) + \lambda_1 \overline{\lambda_2} A_2 \left(\sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} - 1 \right) \right\} \\ B_2(\lambda) &= \lambda_2 \cdot 1_{\mathcal{H}} - \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} (1_{\mathcal{H}} - \overline{\lambda_1} A_1 - \overline{\lambda_2} A_2)^{-1} \\ &\quad \times \left\{ \overline{\lambda_1} \lambda_2 A_1 \left(\sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} - 1 \right) + A_2 \left(|\lambda_1|^2 + |\lambda_2|^2 \sqrt{1 - |\lambda_2|^2 - |\lambda_1|^2} \right) \right\}. \end{aligned}$$

Formulas (4.3), (4.4), and (4.5) allow us to find the explicit forms of operators $\Phi_\lambda(A)$, Ω and Ω_* . On the other hand, from (4.2) follows that if D_{A^*} is one to one, then $D_{\Phi_\lambda(A)^*}$ is also one to one. Using theorem 1, one can obtain the following characterization for Taylor spectrum of $\Phi_\lambda(A)$ in terms of solutions of equations

$$\theta_A(z_1, z_2) D_A(X) = 0 \quad \text{and} \quad (\theta_A(z_1, z_2))^* D_{A^*}(y) = 0.$$

Proposition 5. $\Phi_\lambda(z_1, z_2) \in \sigma_T^{(1)}(\Phi_\lambda(A))$ if and only if equation $\theta_A(z_1, z_2) D_A(X) = 0$ admits at least two nontrivial solutions X_1 and X_2 such that,

$$X_1 = (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda \begin{pmatrix} y \\ 0 \end{pmatrix}, \quad X_2 = (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda \begin{pmatrix} 0 \\ y \end{pmatrix}, \quad y \in \mathcal{H}.$$

Proof. According to Proposition 3, $\Phi_\lambda(z_1, z_2) \in \sigma_T^{(1)}(\Phi_\lambda(A))$ if and only if there exists a nonnul vector $y \in \mathcal{H}$ such that,

$$Y_1 = \begin{pmatrix} y \\ 0 \end{pmatrix}, \quad Y_2 = \begin{pmatrix} 0 \\ y \end{pmatrix}$$

are solutions of equation

$$\theta_{\Phi_\lambda(A)}(\Phi_\lambda(z_1, z_2)) D_{\Phi_\lambda(A)}(Y) = 0.$$

Using theorem 1 and the fact that Φ_λ is involutive, it is equivalent to the existence of a nonnul vector $y \in \mathcal{H}$ such that,

$$Y_1 = \begin{pmatrix} y \\ 0 \end{pmatrix}, \quad Y_2 = \begin{pmatrix} 0 \\ y \end{pmatrix}$$

are solutions of equation

$$\Omega_*^{-1} \theta_A(z_1, z_2) \Omega D_{\Phi_\lambda(A)}(Y) = \Omega_*^{-1} \theta_A(z_1, z_2) D_A \left((1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda(Y) \right) = 0$$

which is equivalent to the equation

$$\theta_A(z_1, z_2) D_A \left((1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda(Y) \right) = 0.$$

□

Corollary 1. Let $(z_1, z_2) \in \mathbb{C}^2$. Then, $(z_1, z_2) \in \sigma_T^{(1)}(A)$ and $\Phi_\lambda(z_1, z_2) \in \sigma_T^{(1)}(\Phi_\lambda(A))$ if and only if there exists two nonnul vectors x and y in $\mathcal{H}$ such that vectors:

$$X_1 = \begin{pmatrix} x \\ 0 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 0 \\ x \end{pmatrix},$$

$$Y_1 = (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda \begin{pmatrix} y \\ 0 \end{pmatrix} \quad \text{and} \quad Y_2 = (1_{\mathcal{H}} - \Lambda^* A)^{-1} D_\Lambda \begin{pmatrix} 0 \\ y \end{pmatrix}$$

are both solutions of equation

$$\theta_A(z_1, z_2) D_A(X) = 0.$$

Proposition 6. Let $(z_1, z_2) \in \mathbb{C}^2$. Assume that equation $(\theta_A(z_1, z_2))^* D_{A^*}(y) = 0$ admits at least one non trivial solution. Then,

$$\Phi_\lambda(z_1, z_2) \in \sigma_T^{(3)}(\Phi_\lambda(A)).$$

Proof. Note at first that operator D_{Λ^*} is invertible. Since Φ_λ is involutive then, according theorem 1,

$$\begin{aligned}
(\theta_A(z_1, z_2))^* D_{A^*}(y) &= 0 \\
&\Leftrightarrow (\theta_A(\Phi_\lambda(\Phi_\lambda(z_1, z_2))))^* D_{A^*}(y) = 0 \\
&\Leftrightarrow (\Omega_* \theta_{\Phi_\lambda(A)}(\Phi_\lambda(z_1, z_2)) \Omega^{-1})^* D_{A^*}(y) = 0 \\
&\Leftrightarrow \Omega(\theta_{\Phi_\lambda(A)} \Phi_\lambda(z_1, z_2))^* \Omega_*^{-1} D_{A^*}(y) = 0 \\
&\Leftrightarrow (\theta_{\Phi_\lambda(A)} \Phi_\lambda(z_1, z_2))^* \Omega_*^{-1} D_{A^*}(y) = 0 \\
&\Leftrightarrow (\theta_{\Phi_\lambda(A)} \Phi_\lambda(z_1, z_2))^* \Omega_*^{-1} D_{A^*} \left((1_{\mathcal{H}} - \Lambda A^*)^{-1} D_{\Lambda^*} D_{\Lambda^*}^{-1} (1_{\mathcal{H}} - \Lambda A^*) y \right) = 0 \\
&\Leftrightarrow (\theta_{\Phi_\lambda(A)} \Phi_\lambda(z_1, z_2))^* D_{\Phi_\lambda(A)^*} (D_{\Lambda^*}^{-1} (1_{\mathcal{H}} - \Lambda A^*) y) = 0 \\
&\Leftrightarrow (\theta_{\Phi_\lambda(A)} \Phi_\lambda(z_1, z_2))^* D_{\Phi_\lambda(A)^*}(X) = 0
\end{aligned}$$

where

$$X = D_{\Lambda^*}^{-1} (1_{\mathcal{H}} - \Lambda A^*) y.$$

Since y is nonnul then, $X = D_{\Lambda^*}^{-1} (1_{\mathcal{H}} - \Lambda A^*) y$ is also nonnul and according proposition 3, it follows that

$$\Phi_\lambda(z_1, z_2) \in \sigma_T^{(3)}(\Phi_\lambda(A_1, A_2)).$$

□

Proposition 7. $\Phi_\lambda(z_1, z_2) \in \sigma_T^{(2)}(A_1, A_2)$ if and only if the equation $\theta_A(z_1, z_2) D_A(X) = 0$ admits at least one solution X such that

$$X \neq \begin{pmatrix} B_1(\lambda)h - w_1.h \\ B_2(\lambda)h - w_2.h \end{pmatrix}, \quad \forall h \in \mathcal{H}$$

where $(w_1, w_2) = \Phi_\lambda(z_1, z_2)$.

Proof. It follows immediately from proposition 2.

□

REFERENCES

- [1] W. Arveson, Subalgebras of C^* -algebras. III. Multivariable operator theory. Acta Math. 181 (1998), 159-228.
- [2] C. Benhida, D. Timotin, Characteristic functions for multicontractions and automorphisms of the unit ball, Preprint.
- [3] T. Bhattacharyya, J. Eschmeier and J. Sarkar, Characteristic function of a pure contractive tuple, Integral Equations Operator theory. Vol. 53, N°1 (2005), 23-32.
- [4] T. Bhattacharyya, J. Eschmeier and J. Sarkar, On completely non coisometric tuples and their characteristic functions, Preprint.
- [5] R. E. Curto and G. C. Hernandez, Ajoint spectral characterisation of primeness for C^* -algebras. Proceeding of Amer Math Soc., Vol.125, N°11 (1997), 3299-3301.
- [6] R. Curto, Application of several complex variables to multiparameter spectral theory (J. B. Conway and B. B. Morell, eds.), Surveys of Some recent Results in Operator Theory, Vol. II, Longman Publ. co., London (1988), 25-90.
- [7] J. Giménez, Joint spectrum of subnormal n -tuples of composition operators. Proceeding of Amer. Math. Soc., Vol. 130, N° 7, (1669), 2015-2023.
- [8] W. Rudin, Function Theory in the unit ball of C^n . Springer-Verlag, New York-Berlin, 1980.
- [9] B. Sz. Nagy and C. Foias, Harmonic analysis of operators on Hilbert space, North-Holland, 1970.

- [10] J. L. Taylor, A joint spectrum for several commuting operators, J. Funct. Anal., 6 (1970) 72-191.

E-mail address: `bbendoukha@gmail.com`

UNIVERSITY OF MOSTAGANEM, DEPARTMENT OF MATHEMATICS, B.O. 227, MOSTAGANEM (27000),
ALGERIA